

SIS epidemics with household structure: the self-consistent field method

G. Ghoshal, L. M. Sander, *

Michigan Center for Theoretical Physics, Department of Physics, University of Michigan, Ann Arbor, Michigan - 48109, U.S.A.

I. M. Sokolov

Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, 12489 Berlin, Germany

Abstract

We consider a stochastic SIS infection model for a population partitioned into m households assuming random mixing. We solve the model in the limit $m \rightarrow \infty$ by using the self-consistent field method of statistical physics. We derive a number of explicit results, and give numerical illustrations. We then do numerical simulations of the model for finite m and without random mixing. We find in many of these cases that the self-consistent field method is a very good approximation.

Key words: SIS epidemic, Households, Stochastic models, Statistical physics

1 Introduction

In this paper we will give a treatment of an SIS infection model in a population partitioned into households. Our main emphasis will be the use of a method borrowed from statistical physics, the self-consistent field (SCF) technique.

The susceptible-infected-susceptible (SIS) model is one of the basic themes in mathematical epidemiology [1,2]. In this idealization of the spread and persistence of an infection, N individuals are initially either susceptible, S , or infected, I . Each one of the infected is coupled to a certain number of other agents. Each I infects all of the S with which it has contact with a rate B ,

* Corresponding author. email lsander@umich.edu

and each I recovers with rate γ , and immediately becomes susceptible again. Schematically, we can write:



Clearly, at long times the system might achieve an equilibrium where the number of infected fluctuates around a mean. We characterize this state by denoting the mean fraction of population which is infected by f , i.e.,

$$f = \langle I \rangle_t / N, \quad (2)$$

where $\langle \rangle_t$ denotes a time average. We refer to this quantity as the endemic level.

The model is defined by the two parameters, B, γ , and by the contact structure of the population. Two cases of the latter have been studied in some detail. In the first, each individual is equally coupled to all the others in a large population. This random mixing model is rather easy to treat, and a large number of results are known [2]. For example, in the deterministic limit, I obeys:

$$\begin{aligned} dI/dt &= bIS/N - \gamma I \\ S &= N - I \end{aligned} \quad (3)$$

where we have redefined $B = b/N$. It is easy to show that for $b > b_c \equiv \gamma$, and for any initial condition with $I > 0$, the epidemic goes to the endemic level, $f = 1 - \gamma/b$. For $b < b_c$ the infection dies out.

In the second well-studied case the individuals are arranged in some geometric way and couple only to their nearest neighbors. This is called the contact process [3]. In the last section of this paper we give numerical results on a model which interpolates between the two cases.

Ball [4] introduced the SIS household model, which puts more structure into the random mixing case by having two levels of mixing. In this idealization the population, N , is partitioned into m households with n members. The infection rate within the household, w , is different from that with individuals outside, B . This followed work on SIR models with similar structure [5]. Ball's results are for the case $m \rightarrow \infty$ with fixed n , that is, a large population of small households, which is clearly of practical interest.

In this paper we extend Ball's work from a different point of view inspired by techniques in statistical physics. These will be explained in the next section. Briefly, the idea is that for random mixing and a large population, we can

consider that each household is acted on by an average force of infection from outside. This is quite similar to methods used in the theory of magnetism [6], inspired by the work of P. Weiss. Weiss introduced the ‘molecular field’, which is proportional to the mean orientation of the neighbors of a given magnetic ion. The orientation of the ion itself is then determined, and the unknown molecular field is gotten by demanding that the neighbors have the same orientation. (Note that this step assumes that there is no correlation between neighbors.) This step is known as self-consistency, hence the name, self-consistent field. As we will see, f plays the role of the self-consistent field. The technique itself is called self-consistent field (or sometimes mean-field) theory.

Some of our results were already obtained in [4], e.g., the expression for b_c for households of size n , Eq. (27), below. Other explicit results are new. We think that one of the main values of this paper is that this point of view allows us to look at the solutions in a new way, and gives considerable insight.

In the last section we continue the analogy to magnetic systems by doing numerical simulations for finite m , and for systems where the random-mixing assumption is not valid. We find, as in magnetism, that the SCF assumption is remarkably accurate in situations where we would expect it to fail.

2 Stochastic formulation and steady-state solutions

Consider an SIS epidemic in a population partitioned into m households each consisting of n individuals. We suppose that within a household the infection rate per contact is w and between different households B . We write $B = b/[(m-1)n]$. (In the notation of Ball [4], $\lambda_w/n = w$, $\lambda_b = b$). We suppose that b remains finite in the limit $m \rightarrow \infty$. The recovery rate is γ , and a recovered individual can be reinfected at once.

2.1 Self-consistent field equations

We begin intuitively by considering the case $n = 2$ and try to formulate the Chapman-Kolmogorov forward equations for the probability, P_k , to have k infected in a typical household:

$$\begin{aligned}\frac{dP_0}{dt} &= \gamma P_1 - 2bfP_0 \\ \frac{dP_1}{dt} &= 2bfP_0^j - (\gamma + w + bf)P_1 + 2\gamma P_2^j\end{aligned}$$

$$\frac{dP_2}{dt} = (w + bf)P_1^j - 2\gamma P_2 \quad (4)$$

$$f = (P_1 + 2P_2)/2 \quad (5)$$

The terms in b are the mean force of infection between households, and f has the interpretation of the fraction of a typical household that is infected. The point is that the only communication between households is by infection, and for a large number of such contacts, only the average number of infected in the other households is important. In effect, we have assumed that the households are independent. Note that $\sum_k P_k = 1$, and $0 \leq f \leq 1$.

The strategy for solving these equations (in the steady state) is to find the P_k with f as a parameter, and solve for f self-consistently, using Eq. (5), at the end of the calculation. This is a heuristically appealing method. For magnetic models, the approximation is exact in the case of infinite-range interactions (in our case, all households interact equally with all others). Even for short-range interactions (i.e., households interacting with a ‘neighborhood’) the approximation is often surprisingly useful.

In fact, a rigorous proof of the validity of Eqs. (4, 5) is available for this case as well. This is given by Ball [4] who uses the work of Kurtz [7] to prove the following: let $X_k^{(m)}$ be the number of households in the population with exactly k infected. Define $P_k = \lim_{m \rightarrow \infty} X_k^{(m)}/m$. Then P_k satisfies Eqs. (4, 5) for any initial condition with finite density, i.e., with $P_k(0), k > 0$ not all equal to 0. In the last section we will investigate numerically how good this approximation is for finite m and for finite range of interaction.

We note, for later use, that we can give another interpretation to Eq. (4). Suppose we consider a single household with 2 individuals subject to a *fixed* external force of infection. Then we can think of P_k as the probability to have k infected. From standard results on stochastic processes, there is a unique steady-state solution to these equations for any positive f, b, w . Further, if $f = 0$ the only steady-state solution is $P_0 = 1, P_1 = P_2 = 0$; that is, the infection becomes extinct with probability unity.

In the general case the time evolution of the vector of probabilities $\mathbf{P} = (P_0, P_1, \dots, P_n)^T$ (T denotes a column vector) to have exactly k infected in the household of n , is given by:

$$\frac{d\mathbf{P}}{dt} = \mathbf{M}(f)\mathbf{P} \quad (6)$$

where $\mathbf{M}(f)$ is a square tridiagonal matrix with nonzero elements

$$m_{k,k-1} = (n - k + 1) [(k - 1)w + fb] \quad (7)$$

$$\begin{aligned} m_{k,k} &= -[k\gamma + (n-k)kw + (n-k)fb] \\ m_{k,k+1} &= (k+1)\gamma \end{aligned}$$

($0 \leq k \leq n$). The equations again contain f , the mean level of infection, as a parameter.

$$f = \frac{1}{n} \sum_{k=0}^n kP_k \quad (8)$$

For the reasons given above, these equations are valid in the limit $m \rightarrow \infty$.

We will be interested in the stationary solutions of Eqs. (6, 8). It is clear by inspection that there is always a trivial steady-state solution $f = 0, P_0 = 1, P_k = 0, k > 0$ corresponding to extinction of the infection. We will find that above a certain threshold there is another solution, and we will show that when this occurs the trivial solution is unstable.

2.2 Steady-state solutions

In the case $n = 2$ it is a matter of simple algebra to find the steady-state solution:

$$\begin{aligned} P_1 &= \frac{2fb\gamma}{\gamma^2 + 2fb\gamma + (fb)^2 + fbw} \\ P_2 &= \frac{(fb)^2 + fbw}{\gamma^2 + 2fb\gamma + (fb)^2 + fbw} \end{aligned} \quad (9)$$

Using Eq. (5) we get a cubic polynomial equation to solve for the endemic level, f . After removing the trivial solution we have:

$$(fb)^2 + f(2b\gamma + wb - b^2) + (\gamma^2 - b\gamma - wb) = 0 \quad (10)$$

The larger root of this equation is:

$$f = \frac{(b - w - 2\gamma) + \sqrt{(b + w)^2 + 4w\gamma}}{2b} \quad (11)$$

This expression is acceptable, i.e., positive, provided:

$$b(\gamma + w)/\gamma^2 > 1 \quad (12)$$

Thus Eq. (12) serves as a threshold condition for the existence of a non-trivial endemic level.

For $n = 3$ it is not difficult to find a cubic equation for f analogous to Eq. (10) (in this expression we set $\gamma = 1$, which sets the unit of time):

$$f^3 b^3 - f^2(b^3 - 3wb^2 - 3b^2) - f(b(2b + 3bw) - 2bw^2 - 3b - 3bw) - b(1 + 2w + 2w^2) + 1 = g(f) = 0 \quad (13)$$

By inspection $g(1) > 0$. If $g(0) = -b(1 + 2w + 2w^2) + 1 < 0$ then there is at least one root in the interval $[0, 1]$. (In fact, numerics shows that there are also two negative roots). Thus if:

$$b(1 + 2w + 2w^2) > 1 \quad (14)$$

we have a non-trivial solution, and otherwise all the roots of Eq. (13) are all negative, and only the trivial solution is acceptable.

In the general case we look for the stationary solution of Eq. (6) by setting $d\mathbf{P}/dt = 0$. We get a homogeneous system of equations, from which the endemic level has to be established self-consistently. Recall that $\sum_k P_k = 1$, so that we can reduce the number of equations by one by eliminating P_0 . Denote the vector of length n , $\mathbf{p} = (P_1, \dots, P_n)^T$. The resulting system of equations for the stationary state is inhomogeneous. It reads

$$\begin{aligned} \mathbf{S}(f)\mathbf{p} &= -nfb\mathbf{u} \\ \mathbf{u} &= (1, 0, 0, \dots)^T \end{aligned} \quad (15)$$

The matrix $\mathbf{S}$ is given by:

$$\begin{pmatrix} m_{1,1} - nfb & m_{1,2} - nfb & -nfb & -nfb & \dots & -nfb \\ m_{21} & m_{22} & m_{23} & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & m_{k,k-1} & m_{k,k} & m_{k,k+1} & 0\dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 & m_{n,n-1} & m_{nn} \end{pmatrix} \quad (16)$$

The matrix $\mathbf{S}(f)$ is linear in f :

$$\mathbf{S}(f) = \mathbf{W} + fb\mathbf{B} \quad (17)$$

where both matrices, $\mathbf{W}$ and $\mathbf{B}$ are f -independent. The structure of $\mathbf{W}$ is:

$$\begin{pmatrix} -\gamma - (n-1)w & 2\gamma & 0 & \dots \\ (n-1)w & -2\gamma - 2(n-2)w & 3\gamma & \dots \\ 0 & 2(n-2)w & -3\gamma - 3(n-3)w & 4\gamma \\ 0 & 0 & 3(n-3)w & \dots \\ 0 & \dots & (n-1)w & -n\gamma \end{pmatrix} \quad (18)$$

$\mathbf{W}$ depends only on the parameters γ and w , characterizing the infection within a household. Note that it is nondegenerate: $\det \mathbf{W} = (-\gamma)^n n!$. To see this, add to each row the sum of all the rows below. The result is a lower triangular matrix with diagonal elements in the n -th row equal to $-n\gamma$. Moreover, $\mathbf{B}$ is a matrix with integer components:

$$\mathbf{B} = \begin{pmatrix} -(2n-1) & -n & -n & -n & \dots \\ (n-1) & -(n-2) & 0 & 0 & \dots \\ 0 & (n-2) & -(n-3) & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & 0 \end{pmatrix}. \quad (19)$$

We give the explicit forms of the matrices $\mathbf{W}$ and $\mathbf{B}$ for $n = 2$ and $n = 3$, setting $\gamma = 1$: For $n = 2$ one has

$$\mathbf{W} = \begin{pmatrix} -1-w & 2 \\ w & -2 \end{pmatrix} \text{ and } \mathbf{B} = \begin{pmatrix} -3 & -2 \\ 1 & 0 \end{pmatrix}, \quad (20)$$

For $n = 3$ we get

$$\mathbf{W} = \begin{pmatrix} -1-2w & 2 & 0 \\ 2w & -2-2w & 3 \\ 0 & 2w & -3 \end{pmatrix} \text{ and } \mathbf{B} = \begin{pmatrix} -5 & -3 & -3 \\ 2 & -1 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \quad (21)$$

From the remarks above, it is clear that there is a non-zero solution, $\mathbf{p}$, to Eq. (15) for any $f > 0$, so that $\mathbf{S}^{-1}$ exists. Thus:

$$\mathbf{p} = -nfb\mathbf{S}^{-1}\mathbf{u} \quad (22)$$

From this equation a closed form for the endemic level f follows. Multiplying both sides of Eq.(22) by the vector $\mathbf{n}_1 = (1, 2, \dots, n)$ and noting that $f = n^{-1}\mathbf{n}_1\mathbf{p}$ we get:

$$f = -f\mathbf{n}_1\mathbf{S}^{-1}\mathbf{u} \quad (23)$$

which is a closed algebraic equation for f which always has the trivial solution $f = 0$. Factoring out this out, we get an equation for the nontrivial solution:

$$-b\mathbf{n}_1\mathbf{S}^{-1}\mathbf{u} = 1. \quad (24)$$

As in Eqs. (12, 14) there is a condition on the parameters for the solution to be positive. For given w we define a value b_c such that for $b > b_c$ we have such an f . This can be found from Eq.(24) by noting that at b_c the second solution crosses $f = 0$. At that point, $\mathbf{S} = \mathbf{W}$, and we get:

$$b_c = -1 / [\mathbf{n}_1\mathbf{W}^{-1}\mathbf{u}]. \quad (25)$$

To get an explicit expression for b_c consider the vector $\mathbf{v} = \mathbf{W}^{-1}\mathbf{u}$, whose elements are $\mathbf{v}_k = [\mathbf{W}^{-1}]_{1,k}$. As we will show,

$$\mathbf{v}_k = -\frac{(n-1)!}{k(n-k)!}w^{k-1}, \quad (26)$$

so that the threshold is given by

$$b_c^{-1} = -\mathbf{n}_1\mathbf{v} = \sum_{k=1}^n \frac{(n-1)!}{(n-k)!}w^{k-1}. \quad (27)$$

To prove Eq.(26) we note that the solution to $\mathbf{W}\mathbf{v} = \mathbf{u}$ is unique so that we need only insert Eq. (26) into $\mathbf{W}\mathbf{v} = \mathbf{u}$. The first element is

$$-(-1 - (n-1)w) - (n-1)w = 1, \quad (28)$$

while the k -th element ($k > 1$) is:

$$\begin{aligned} & (k-1)(n+1-k)w \cdot \frac{(n-1)!}{(n+1-k)!(k-1)!}w^{k-2} \\ & + (-k - k(n-k)w) \frac{(n-1)!}{(n-k)!k}w^{k-1} + \end{aligned}$$

$$\begin{aligned}
& +(k+1) \frac{(n-1)!}{(n-k-1)!(k+1)} w^k = \\
& = (n-1)! \left\{ \left[\frac{1}{(n-k)!} - \frac{1}{(n-k)!} \right] w^{k-1} + \right. \\
& \quad \left. + \left[\frac{1}{(n-k-1)!} - \frac{1}{(n-k-1)!} \right] w^k \right\} \\
& = 0.
\end{aligned} \tag{29}$$

Once we obtain $\mathbf{p}$ we can find other moments of the distribution by multiplying Eq.(22) by $\mathbf{n}_q = (1, 2^q, 3^q, \dots, n^q)$. We present numerical results on $\mathbf{n}_2\mathbf{p}$ below.

We now discuss the behavior of the solution in the vicinity of the transition point. Eq.(22) can be rewritten:

$$\mathbf{p} = -nfb \left(\mathbf{I} + fb\mathbf{W}^{-1}\mathbf{B} \right)^{-1} \mathbf{W}^{-1}\mathbf{u}, \tag{30}$$

Then we formally expand in a geometric series:

$$\mathbf{p} = -nfb \left(\mathbf{I} - fb\mathbf{W}^{-1}\mathbf{B} + (fb)^2 \left(\mathbf{W}^{-1}\mathbf{B} \right)^2 - \dots \right) \mathbf{W}^{-1}\mathbf{u}. \tag{31}$$

Multiplying both sides of Eq.(31) by $n^{-1}\mathbf{n}_1$ we get the following equation for f :

$$f = -fb \left(a_0 - fba_1 + (fb)^2 a_2 + \dots + (-1)^m (fb)^m a_m + \dots \right) \tag{32}$$

where

$$a_m = \mathbf{n}_1 \left(\mathbf{W}^{-1}\mathbf{B} \right)^m \mathbf{W}^{-1}\mathbf{u} \tag{33}$$

from which f can be obtained with any necessary accuracy. Note that according to Eq.(25) the coefficient

$$a_0 = \mathbf{n}_1 \mathbf{W}^{-1}\mathbf{u} = -1/b_c. \tag{34}$$

Eq.(32) is especially useful in the vicinity of transition, where f is small. Both the point of transition and the endemic level near the threshold can be determined.

Eq.(32) always possesses a trivial solution $f = 0$. An equation for a nontrivial one then reads:

$$G(f) = fb^2 a_1 - f^2 b^3 a_2 + \dots + (-1)^{m+1} f^m b^{m+1} a_m + \dots = 1 + ba_0. \tag{35}$$

The formal solution of this equation involves the inverse of G :

$$f = G^{-1}(1 + ba_0) = G^{-1}(1 - b/b_c). \quad (36)$$

where Eq (34) has been used. Using the relationship between the Taylor series of a function and its inverse we get, in the vicinity of b_c :

$$f = -\beta_1\tau + \beta_2\tau^2 - \dots \quad (37)$$

where $\tau = (b - b_c)/b_c$ and the coefficients are:

$$\begin{aligned} \beta_1 &= \frac{1}{b^2 a_1(w)}, \\ \beta_2 &= -\frac{a_2(w)}{b^3 a_1^3(w)}, \\ \beta_3 &= \frac{1}{b^4 a_1^5(w)} (2a_2(w) - a_1(w)a_3(w)), \\ &\dots \end{aligned} \quad (38)$$

In the vicinity of the transition one can take $b = b_c(w)$; the behavior of f is dominated by the linear term:

$$f = -\beta_1\tau = -\frac{1}{b^2 a_1(w)}\tau \quad (39)$$

Explicit calculations show that $a_1 < 0$.

2.3 Stability of the disease-free equilibrium

Restoring the time-dependence in Eq.(15) we obtain an equation for the time evolution of the reduced probability vector $\mathbf{p}$:

$$\frac{d}{dt}\mathbf{p} = (\mathbf{W} + fb\mathbf{B})\mathbf{p} + nfb\mathbf{u}. \quad (40)$$

Since $\mathbf{p}$ depends on f , Eq.(40) is a nonlinear equation describing the evolution of $\mathbf{p}$. In the previous section we have shown that this equation has two equilibria provided $b > b_c$, where b_c is defined in Eq. (27). We will now show that under the same conditions the trivial solution, $f = 0, \mathbf{p} = \mathbf{0}$ is unstable.

In the vicinity of the trivial equilibrium, we can omit the term $fb\mathbf{B}$ in brackets in Eq. (40). The linearized equation reads

$$\frac{d}{dt}\mathbf{p} = \mathbf{W}\mathbf{p} + b(\mathbf{n}_1\mathbf{p})\mathbf{u} \equiv \mathbf{T}\mathbf{p} \quad (41)$$

where

$$\mathbf{T} = \mathbf{W} + b\mathbf{N}, \quad (42)$$

The matrix $\mathbf{N}$ has $\mathbf{n}_1$ as its first row and all other elements zero.

The determinant of $\mathbf{T}$ is linear in b . To see this, note that the explicit form of $\mathbf{T}$ is:

$$\begin{pmatrix} -\gamma - (n-1)w + b & 2\gamma + 2b & 3b & \dots \\ (n-1)w & -2\gamma - 2(n-2)w & 3\gamma & \dots \\ 0 & 2(n-2)w & -3\gamma - 3(n-3)w & \dots \\ \dots & \dots & \dots & \dots \\ 0 & \dots & (n-1)w & -n\gamma \end{pmatrix} \quad (43)$$

If we expand the determinant about the first row, each term contains b only once. It can thus be written $\det \mathbf{T} = \det \mathbf{W} + cb$, with c a constant. However, the determinant of $\mathbf{T}$ is the product of its eigenvalues. Since for $b = 0$ (separate households) the trivial solution is stable, all of the eigenvalues of $\mathbf{W}$ are negative; $\det \mathbf{W}$ is nonzero, and its sign is positive for n even and negative for n odd. From the linearity in b we see that $\det \mathbf{T}$ changes sign at $b = -c^{-1} \det \mathbf{W}$ which corresponds to the appearance of a positive eigenvalue.

To find the b for which $\det \mathbf{T} = 0$ write Eq.(42) as:

$$\mathbf{T}\mathbf{W}^{-1} = \mathbf{I} + b\mathbf{N}\mathbf{W}^{-1}. \quad (44)$$

The determinant of the matrix in the l.h.s. of this expression is

$$\det \mathbf{T} / \det \mathbf{W} = [(-\gamma)^{-n} / n!] \det \mathbf{T} \quad (45)$$




Fig. 1. The endemic level as a function of b for $n = 3$ and various w , $\gamma = 1$

and vanishes when $\det \mathbf{T} = 0$. But, since $\mathbf{N}$ has non-zero elements only in the first row, the matrix on the r.h.s. of Eq. (44) is:

$$\begin{pmatrix} 1 + bq_1 & bq_2 & bq_3 & \dots & bq_n \\ 0 & 1 & 0 & \dots & \dots \\ 0 & 0 & 1 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 & 1 \end{pmatrix}, \quad (46)$$

where q_i is the i -th element of the vector $\mathbf{n}_1 \mathbf{W}^{-1}$. By inspection, the determinant of this matrix is $1 + q_1 \equiv 1 + b \mathbf{n}_1 \mathbf{W}^{-1} \mathbf{u}$. It vanishes at $b = -1/(\mathbf{n}_1 \mathbf{W}^{-1} \mathbf{u})$, i.e. exactly at $b = b_c$, see Eq.(25). Therefore, for $b > b_c$ a positive eigenvalue appears, and the trivial solution loses its linear stability.

2.4 Numerical results

It is quite simple to work out the dependence of f on the parameters w, b, n . We wrote a Matlab program which solved Eq. (24). In Figure (1) we show f




Fig. 2. The standard deviation of the number of infected in a household, as a function of b for $n = 3$ for several w 's, $\gamma = 1$




Fig. 3. The endemic level as a function of b for $w = 0.2$ and $n = 1, 2, \dots, 8$, $\gamma = 1$




Fig. 4. The standard deviation of the number of infected in a household, as a function of b for $w = 0.2$ and $n = 1, 2, \dots, 8$, $\gamma = 1$

as a function of b for several w 's for $n = 3$. In this and the following figures we have taken $\gamma = 1$. For other values of γ one can use the obvious scaling relation $(w, b, \gamma) \rightarrow (w/\gamma, b/\gamma, 1)$.

Recall that $f = \mathbf{n}_1 \mathbf{p}$ can be interpreted as $\langle i \rangle / n$ where i is the number infected in a household, and $\langle \rangle$ is the average over the distribution $\mathbf{P}$. If we calculate $\mathbf{n}_2 \mathbf{p}$ we get a measure of the fluctuation of i in a household. In Figure (2) we show the standard deviation, $\sigma = [\langle i^2 \rangle - \langle i \rangle^2]^{1/2} = [\mathbf{n}_2 \mathbf{p} - (nf)^2]^{1/2}$ as a function of b for several w 's, for $n = 3$. In a similar way, in Figures (3, 4) we show f and σ for $w = 0.2$ and $n = 1, 2, \dots, 8$.

3 Simulations: finite systems and finite range

The considerations in the previous sections apply in the limit $m \rightarrow \infty$. For any finite m the only steady-state solution possible is the trivial one, $f = 0$; with probability unity the infection will become extinct. However, the time to extinction may be very long, $O(e^m)$ if the system is above b_c . See, for example, [8,9,10] for the case $n = 1$. For times short compared to the extinction time the system will be in a quasi-stationary state which resembles the infinite system.




Fig. 5. The coupling scheme used in the simulations. This is the case $n = 3$, $m = 8$, $L = 2$. The white bonds represent contacts within a household. All of the inter-household contacts for one agent are shown in black

We have done preliminary investigations of this and related behavior using computer simulations. We represent the population as follows: imagine a population of m households each containing n agents arranged on a ring. All of the members of the same household can infect one another with probability/unit time w . The communication with other households is variable range: we allow each household to infect all of the other households in a range L on either side. See Figure (5). That is, the number of outside contacts is $q = 2Ln$, and the infection probability is b/q . The equivalent of the random mixing case of the previous sections is $2L = m - 1$; in this case all agents are coupled to each other. The case $n = 1$, $L = 1$ is the well-known contact process [3].

Our computer algorithm is quite simple: we suppose that γ, w, b are all less than unity. We start by marking a certain fraction of agents as infected, which establishes $I(0)$ where I is the total number infected in the population. For each computer time step we choose one agent, in household k , at random among the infected, and find all the other agents that are coupled to the one in question. These will be in households with index $k - L, k - L + 1, \dots, k + L$, where the indices are to be interpreted with periodic boundary conditions. With probability w or b/q (depending whether the other agent is in k or outside) the other agent is infected. Then another agent is chosen at random among the infected, and it is allowed to recover with probability γ . Each computer time step corresponds to an increment of ‘clock time’, $dt = 1/I$. A typical result for $I(t)$ is shown in Figure(6). It is worth noting that this algorithm is quite fast – it is not difficult to treat as many as 3000 agents on a simple workstation.

In this section we will give a few results of the simulations. We have found




Fig. 6. Simulation results for the number of infected as a function of time compared to the SCF results. $n = 3$, $m = 501$, $L = 150$, $w/\gamma = 0.2$, $b/\gamma = 1$. The agreement for the average is essentially perfect, but for the fluctuations (see text) the SCF method is an underestimate.

that the SCF is an excellent representation of the behavior in large systems with long-range interactions, but that for smaller systems and shorter ranges the approximation becomes poorer, but still surprisingly good. A systematic investigation of these effects will be reserved for a future publication: here we give a few preliminary results.

3.1 Finite size effects

For large enough m the simulation results agree very well with the SCF results. In Figure(6) the average of the time series agrees almost perfectly with the results from the previous sections: the numerical average of the simulation data differs from the SCF by less than the width of the line.

In addition we can try to understand the fluctuations of the time series by the

following reasoning: if we imagine that the different households fluctuate more or less independently (this is the assumption under which the SCF method is valid), then the standard deviation of the total should be $\sigma_{tot} = \sigma\sqrt{m}$. As we see in the figure, this is a good approximation. However, we have observed that there is a large finite-size effect for this statistic. For $m = 301$ the simulation result for σ_{tot} is almost twice the prediction.

Another way to look at the standard deviation is to simply record the time series of the number infected in a given household, and directly find σ . We did this by averaging over 10 randomly chosen households in the system. We find very good agreement with the analytic treatment: for example, for $w = 0.2, b = 1, n = 3, m = 501$ we find $\sigma = 0.82$. The analytic result is 0.84, well within our estimated error.

As we reduce m we find a surprising result: within the numerical accuracy of our simulation we can find no change in f . Rather, at a certain small size, $m = 50$ for the parameters in the last paragraph, the infection dies before we can collect adequate statistics.

3.2 Finite range effects

For the case of small L the SCF ceases to be valid. In magnetic systems, locality of the interaction is well known to increase fluctuations and to change the threshold. We find related effects here, but the range needs to be very small indeed for the SCF result to be very inaccurate.

In Figure (7) we show simulation results for $f(L)$, the endemic level as a function of the range of the interaction between households, L . We show three different sets of parameters for $n = 3$ and compared with the SCF ($L \rightarrow \infty$). The dependence is remarkable: for the parameter values ($w = 1, b = 1$), finite range has hardly any effect. The other two sets do show a decrease of f , and an apparent shift in the threshold. For $w = 0.2, b_c = 0.67$ in the SCF, well below $b = 1$. However, for $L < 5$, the system is below threshold, and the epidemic dies quickly. A similar effect occurs for $w = 1, b = 0.3$, which is below b_c for $L < 10$. (For large L , $b_c = 0.2$ for these parameters.) Another way to put this is that there is a characteristic length $L_c \approx 10$ for these two sets of parameters.

These results are reminiscent of effects that occur in phase transitions in statistical physics. In fact, the contact process is fruitfully viewed as an example of a continuous non-equilibrium phase transition [11]. Near the threshold (i.e. the critical point) of a continuous phase transition, the system develops a diverging correlation length. For total system sizes greater than this length finite size effects are small. Thus we might guess that the correlation lengths for two of the sets of parameters could be of order L_c . We have not investigated this




Fig. 7. The endemic level, f , as a function of the range of the infection contacts between households. The horizontal lines are the SCF results for large L . In all cases $n = 3, m = 501$ so that the maximum possible $L = 250$

matter in detail, but it would repay consideration.

4 Summary

In this paper we have given a number of explicit results for the SIS household model. Our point of view was to emphasize the SCF method as a guide to solution and insight. The numerical results give rise to a number of questions that are probably not accessible to rigorous proof, but for which numerical methods could be pursued. For example, we have speculated that there are critical fluctuations [6,11] in this system. That would mean that near threshold there would be large correlations in adjacent households for a finite range system, a result that could be of considerable interest.

There is also a practical aspect to this method. SCF theory is quite flexible, and allows considerable complication to be added to the model without sub-

stantially increasing the complexity of the solution. Suppose our household contains two types of members, for example, children who go to school and contract infections, and adults who do not. This would require a few changes: for a 3 person household with two adults and a child, the analog of the vector $\mathbf{P}$ would be of length 6 instead of 4, and the complete solution would be no worse than the numerical inversion of a 5×5 matrix.

Acknowledgements

We would like to thank Frank Ball, Mark Newman, Carl Simon, and Jim Koopman for helpful discussions. GG was supported by the REU summer program of the Physics Department of the University of Michigan, and IMS would like to thank the Michigan Complex Systems program for support.

References

- [1] R. Ross, Some a priori pathometric equations, *Br. Med. J.* 1 (1915) 895.
- [2] H. W. Hethcote, Qualitative analysis of communicable disease models, *Math. Biosci.* 28 (1976) 335.
- [3] T. M. Liggett, *Interacting Particle Systems*, Springer, 1985
- [4] F. Ball, Stochastic and deterministic models for SIS epidemics among a population partitioned into households, *Math. Biosci.*, 156 (1999) 41.
- [5] F. G. Ball, D. Mollison, G. Scalia-Tomba, Epidemics with two levels of mixing, *Ann. Appl. Prob.* 7 (1997) 46.
- [6] K. Huang, *Statistical Mechanics*, Wiley, 1987
- [7] T. G. Kurtz, Solutions of ordinary differential equations as limits of pure jump Markov processes, *J. Appl. Prob.* 7 (1970) 49.
- [8] G. H. Weiss and M. Dishon, On the asymptotic behavior of the stochastic and deterministic models of an epidemic, *Math. Biosci.* 11 (1971) 261.
- [9] I. Nasell, The quasi-stationary distribution of a closed endemic SIS model, *Adv. Appl. Prob.* 28 (1996) 895.
- [10] H. Andersson and B. Djehiche, A threshold limit theorem for the stochastic logistic epidemic, *J. Appl. Prob.* 35 (1998) 662.
- [11] J. Marro and R. Dickman, *Nonequilibrium phase transitions in lattice models*, Cambridge University Press, 1999